

Nonlinear Plasma Response

G. ADOMIAN AND R. RACH

Center for Applied Mathematics, University of Georgia, Athens, Georgia 30602

Submitted by E. Stanley Lee

The decomposition method (G. Adomian, "Stochastic Systems," Academic Press, New York/London, 1983; G. Adomian, "Stochastic Systems II," in press) (and earlier work of Adomian referenced therein) allows solution without linearization of problems normally requiring the self-consistent field technique. The latter technique can provide a sufficiently accurate approximation in weakly coupled plasma, but the decomposition method of Adomian provides a superior and physically preferable methodology for the general case. © 1985 Academic Press, Inc.

The response of an interacting electron stream to electrostatic perturbation is widely considered to be best calculated by the self-consistent-field technique [1] developed by Ehrenreich and Cohen [2] and Goldstone and Gottfried [3]. In seeking the equation for the charge density, use of the self-consistent field method requires taking the expectation value of the equation of motion whereas one should solve the equation first as in the decomposition method [4, 5] and then average. The usual method is a typical hierarchy procedure which necessarily neglects interaction terms simply replacing an average of a product by a product of averages. While this can be a sufficiently accurate approximation in weakly coupled plasmas, the error in doing so in general has been well established by Adomian [4, 6], Adomian and Malakian [7]. It is in fact (as shown in [4]) a perturbation method in which the instantaneous field experienced by an electron does not deviate much from the average value. We show here using the methodology of [4, 5] that a better, and certainly more physically realistic method of solution is possible now.

The total charge density consists of two parts—an external charge density ρ_{ext} and an induced charge density ρ_{ind} due to the nonuniformity of the plasma. The electrostatic potential in a single-component plasma can be calculated from Poisson's equation

$$\nabla^2 \phi = -\rho_{\text{ext}} + \alpha(e^{-\beta\phi} - 1) \quad (1)$$

where the second term on the right is the induced charge ρ_{ind} . Thus, we have a nonlinear differential equation for ϕ . At this point it is customary to

make various approximations, e.g., ρ_{ext} is assumed small so that the induced potential will also be small ($\beta\phi \ll 1$) and one can write a linearized equation. However, we make no linearizing assumption. We write (1) in the standard form of [4, 5] as

$$\nabla^2 \phi - \alpha e^{-\beta\phi} = -\rho_{\text{ext}} - \alpha \quad (2)$$

or $F\phi \equiv L\phi + N\phi = g$, where $L = \nabla^2$ and $N\phi = -\alpha e^{-\beta\phi}$ is a nonlinear term, and $g = -\rho_{\text{ext}} - \alpha$. Let $L = L_x + L_y + L_z$, where $L_x = \partial^2/\partial x^2$, $L_y = \partial^2/\partial y^2$, $L_z = \partial^2/\partial z^2$ (see [5, 9]). Then

$$[L_x + L_y + L_z]\phi + N\phi = g.$$

We solve for $L_x\phi$, $L_y\phi$, $L_z\phi$ in turn as discussed in [5, 8] so that

$$L_x\phi = g - L_y\phi - L_z\phi - N\phi$$

$$L_y\phi = g - L_z\phi - L_x\phi - N\phi$$

$$L_z\phi = g - L_x\phi - L_y\phi - N\phi.$$

The inverses L_x^{-1} , L_y^{-1} , L_z^{-1} are defined [5, 8] as two-fold definite integrations (thus L_x is a double integration between the limits 0 to x , etc.) If we indicate the homogeneous solutions by Φ_x , Φ_y , Φ_z (i.e., $L_x\Phi_x = 0$, etc.)

$$\phi = \Phi_x + L_x^{-1}g - L_x^{-1}L_y\phi - L_x^{-1}L_z\phi - L_x^{-1}N\phi$$

$$\phi = \Phi_y + L_y^{-1}g - L_y^{-1}L_z\phi - L_y^{-1}L_x\phi - L_y^{-1}N\phi$$

$$\phi = \Phi_z + L_z^{-1}g - L_z^{-1}L_x\phi - L_z^{-1}L_y\phi - L_z^{-1}N\phi.$$

Adding (and dividing by 3),

$$\begin{aligned} \phi &= \left(\frac{1}{3}\right)\{(\Phi_x + \Phi_y + \Phi_z) + (L_x^{-1} + L_y^{-1} + L_z^{-1})g\} \\ &\quad - \left(\frac{1}{3}\right)\{L_x^{-1}L_y + L_y^{-1}L_z + L_z^{-1}L_x + L_x^{-1}L_z + L_y^{-1}L_x + L_z^{-1}L_y\}\phi \\ &\quad - \frac{1}{3}\{L_x^{-1} + L_y^{-1} + L_z^{-1}\}N\phi. \end{aligned}$$

We define the entire first of the above three lines as ϕ_0 in the decomposition $\phi = \sum_{n=0}^{\infty} \phi_n$. Define the entire second line as $-(L^{-1}R)\phi$ and the third line as $L^{-1}N\phi$ then we have the convenient form

$$\phi = \phi_0 - (L^{-1}R)\phi - L^{-1}(N\phi). \quad (3)$$

For ϕ we substitute $\phi = \sum_{n=0}^{\infty} \phi_n$ and $N\phi$ is developed in Adomian's A_n polynomials [4, 5] thus

$$\sum_{n=0}^{\infty} \phi_n = \phi_0 - (L^{-1}R) \sum_{n=0}^{\infty} \phi_n - L^{-1} \sum_{n=0}^{\infty} A_n. \quad (4)$$

The A_n of course are developed for the particular nonlinearity present [4, 5, 8]. The components of ϕ are

$$\begin{aligned}\phi_0 &= (\tfrac{1}{3})\{(\Phi_x + \Phi_y + \Phi_z) + (L_x^{-1} + L_y^{-1} + L_z^{-1})g\} \\ \phi_1 &= -L^{-1}R\phi_0 - L^{-1}A_0 \\ \phi_2 &= -L^{-1}R\phi_1 - L^{-1}A_1 \\ \phi_3 &= -L^{-1}R\phi_2 - L^{-1}A_2 \\ &\vdots\end{aligned}\tag{5}$$

The A_n have been previously defined (see [4, 5, 8, 9]) for $N\phi = \alpha e^{-\beta\phi}$ they are¹

$$\begin{aligned}A_0 &= \alpha e^{-\beta\phi_0} \\ A_1 &= \alpha e^{-\beta\phi_0}[-\beta\phi_1] \\ A_2 &= \alpha e^{-\beta\phi_0}[-\beta\phi_2 + (\beta^2/2!)\phi_1^2] \\ A_3 &= \alpha e^{-\beta\phi_0}[-\beta\phi_3 + \beta^2\phi_1\phi_2 - (\beta^3/3!)\phi_1^3] \\ A_4 &= \alpha e^{-\beta\phi_0}[-\beta\phi_4 + \beta^2\{(\phi_2^2/2!) + \phi_1\phi_3\} \\ &\quad - (\beta^3\phi_1^2\phi_2/2!) + (\beta^4\phi_1^4/4!)] \\ A_5 &= \alpha e^{-\beta\phi_0}[-\beta\phi_5 + \beta^2\{\phi_2\phi_3 + \phi_1\phi_4\} \\ &\quad - \beta^3\{(\phi_1\phi_2^2/2!) + (\phi_1^2\phi_3/2!)\} \\ &\quad + (\beta^4\phi_1^3\phi_2/3!) - (\beta^5\phi_1^5/5!)] \\ A_6 &= \alpha e^{-\beta\phi_0}[-\beta\phi_6 + \beta^2\{(\phi_3^2/2!) + \phi_2\phi_4 + \phi_1\phi_5\} \\ &\quad - \beta^3\{(\phi_2^3/3!) + \phi_1\phi_2\phi_3 + \phi_1^2\phi_4/2!\} \\ &\quad + \beta^4\{(\phi_1^2/2!)(\phi_2^2/2!) + (\phi_1^3/3!)\phi_3\} \\ &\quad - \beta^5(\phi_1^4/4!)\phi_2 + \beta^6(\phi_1^6/6!)] \\ A_7 &= \alpha e^{-\beta\phi_0}[-\beta\phi_7 + \beta^2\{\phi_3\phi_4 + \phi_2\phi_5 + \phi_1\phi_6\} \\ &\quad - \beta^3\{(\phi_2^2/2!)\phi_3 + \phi_1(\phi_3^2/2!) + \phi_1\phi_2\phi_4 \\ &\quad + (\phi_1^2/2!)\phi_5\} + \beta^4\{\phi_1(\phi_2^3/3!) + (\phi_1^2/2!)\phi_2\phi_3 \\ &\quad + (\phi_1^3/3!)\phi_4\} - \beta^5\{(\phi_1^3/3!)(\phi_2^2/2!) + (\phi_1^4/4!)\phi_3\} \\ &\quad + \beta^6(\phi_1^5/5!)\phi_2 - \beta^7(\phi_1^7/7!)]\end{aligned}\tag{6}$$

¹ We can let $N\phi = e^{-\beta\phi}$ and write αNy instead of Ny .

$$\begin{aligned}
A_8 = & \alpha e^{-\beta\phi_0} [-\beta\phi_8 + \beta^2 \{ (\phi_4^2/2!) + \phi_3\phi_5 + \phi_2\phi_6 + \phi_1\phi_7 \} \\
& - \beta^3 \{ \phi_2(\phi_3^2/2!) + (\phi_2^2/2!) \phi_4 + \phi_1\phi_3\phi_4 \\
& + \phi_1\phi_2\phi_5 + (\phi_1^2/2!) \phi_6 \} \\
& + \beta^4 \{ (\phi_4^4/4!) + \phi_1(\phi_2^2/2!) \phi_3 + (\phi_1^2/2!)(\phi_3^2/2!) \\
& + (\phi_1^2/2!) \phi_2\phi_4 + (\phi_1^3/3!) \phi_5 \} \\
& - \beta^5 \{ (\phi_1^2/2!)(\phi_3^2/3!) + (\phi_1^3/3!) \phi_2\phi_3 + (\phi_1^4/4!) \phi_4 \} \\
& + \beta^6 \{ (\phi_1^4/4!)(\phi_2^2/2!) + (\phi_1^5/5!) \phi_3 \} \\
& - \beta^7 (\phi_1^6/6!) \phi_2 + \beta^8 (\phi_1^8/8!)] \\
A_9 = & \alpha e^{-\beta\phi_0} [-\beta\phi_9 + \beta^2 \{ \phi_4\phi_5 + \phi_3\phi_6 + \phi_2\phi_9 + \phi_1\phi_8 \} \\
& - \beta^3 \{ (\phi_3^3/3!) + \phi_2\phi_3\phi_4 + (\phi_2^2/2!) \phi_5 + \phi_1(\phi_2^2/2!) \\
& + \phi_1\phi_3\phi_5 + \phi_1\phi_2\phi_6 + (\phi_1^2/2!) \phi_7 \} \\
& + \beta^4 \{ (\phi_3^2/3!) \phi_3 + \phi_1\phi_2(\phi_3^2/2!) + \phi_1(\phi_2^2/2!) \phi_4 \\
& + (\phi_1^2/2!) \phi_3\phi_4 + (\phi_1^2/2!) \phi_2\phi_5 + (\phi_1^3/3!) \phi_6 \} \\
& - \beta^5 \{ \phi_1(\phi_2^4/4!) + (\phi_1^2/2!)(\phi_2^2/2!) \phi_3 \\
& + (\phi_1^3/3!)(\phi_3^2/2!) + (\phi_1^3/3!) \phi_2\phi_4 + (\phi_1^4/4!) \phi_5 \} \\
& + \beta^6 \{ (\phi_1^3/3!)(\phi_2^3/3!) + (\phi_1^4/4!) \phi_2\phi_3 + (\phi_1^5/5!) \phi_4 \} \\
& - \beta^7 \{ (\phi_1^5/5!)(\phi_2^2/2!) + (\phi_1^6/6!) \phi_3 \} \\
& + \beta^8 (\phi_1^7/7!) \phi_2 - \beta^9 (\phi_1^9/9!)]
\end{aligned}$$

which is sufficient to give us the 10-term approximation ϕ_{10} . (More are easily calculated if the problem warrants it, but we expect a very rapidly damped oscillating convergence from results discussed previously—particularly in [5].)

Since $\beta\phi$ is generally assumed small ($\beta\phi \ll 1$), $e^{-\beta\phi} - 1$ becomes $-\beta\phi$ to make a linear equation. The decomposition method solves Eq. (1) *without linearizing assumptions, perturbative methods, or truncations* and the complete solution is given by

$$\phi_n = \sum_{v=0}^{n-1} \phi_v$$

with the ϕ_v given by (5) using the A_n in (6).

Complete and accurate solution can now be obtained for any specific initial/boundary conditions for (1) and values of α , β , ρ_{ext} in a rapidly convergent series [10] to any desired approximation.

REFERENCES

1. P. M. PLATZMAN AND P. A. WOLFF, "Waves and Interactions in Solid State Plasmas," Academic Press, New York/London, 1973.
2. H. EHRENREICH AND M. H. COHEN, *Phys. Rev.* **115** (1959), 786.
3. J. GOLDSTONE AND K. GOTTFRIED, *Nuovo Cimento (10)* **13** (1959), 849.
4. G. ADOMIAN, "Stochastic Systems," Academic Press, New York/London, 1983.
5. G. ADOMIAN, "Stochastic Systems II," Academic Press, New York/London, in press.
6. G. ADOMIAN, The closure approximation in the hierarchy equations, *J. Statist. Phys.* **3**, No. 2, (1971).
7. G. ADOMIAN AND K. MALAKIAN. Closure approximation error in the mean solution of stochastic differential equations by the hierarchy method, *J. Statist. Phys.* **21**, No. 2, (1979), 181–189.
8. R. E. BELLMAN AND G. ADOMIAN, "Partial Differential Equations—New Methods for Their Treatment and Applications," Reidel, Dordrecht, 1985.
9. R. RACH, A convenient computational form for the Adomian polynomials, *J. Math. Anal. Appl.* **102**, No. 2, (1984), 415–419.
10. G. ADOMIAN, Convergent series solution of nonlinear equations, *J. Comput. Appl. Math.* **11** (1984), 225–230.